DOI: 10.2174/0125902776419140251028065415, 2025, 18, e25902776419140

RESEARCH ARTICLE OPEN ACCESS

Leaf Litter Decomposition Rate in the Karst Ecosystem of the Community Forest in Wonodadi Village, Pracimantoro, Wonogiri, Indonesia



ISSN: 2590-2776

Rieska Rahayu Safitri¹, Malihatun Nufus^{1,*} and Supriyadi¹

 1 Department of Forest Management, Faculty of Agriculture, Universitas Sebelas Maret, Surakarta, Indonesia

Abstract:

Introduction: The availability of mineral nutrients in the karst ecosystem is specifically influenced by vegetation, which depends on how much nutrition is contributed by the litter of the stand through the decomposition process. The litter decomposition rate can be observed through physical changes and a decrease in litter weight. The community in Pracimantoro utilizes and maintains land by building community forests to optimize land in the karst area. This study aims to determine the litter decomposition rate in several community forest planting patterns in Pracimantoro.

Methods: Data were collected using the litter bag method by making sample plots measuring 0.04 ha and selected by purposive sampling, and placed diagonally. The rate of decomposition is described by the constant value (k).

Results: The karst community forest in Pracimantoro was dominated by Mahogany (Swietenia sp.), Acacia (Acacia auriculiformis), and Teak (Tectona grandis) stands. The rate of decomposition in the Pracimantoro karst community forest was 0.0268.

Discussion: Leaf litter decomposition over two months resulted in a decrease in litter weight of 33.05%. The duration of nutrient return in karst community forests can be estimated using the half-life approach.

Conclusion: The rate of litter decomposition is influenced by environmental conditions and litter quality. Environmental conditions that are suitable over a long period have a significant impact on the rate of litter decomposition.

Keywords: Litter stock, Half-life approach, Associated plant species, Climate variable, Swietenia sp.

© 2025 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: https://creativecommons.org/licenses/by/4.0/legalcode. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Address correspondence to this author at the Department of Forest Management, Faculty of Agriculture, Universitas Sebelas Maret, Surakarta, Indonesia; E-mail: malihatunufus@staff.uns.ac.id

Cite as: Safitri R, Nufus M, Supriyadi . Leaf Litter Decomposition Rate in the Karst Ecosystem of the Community Forest in Wonodadi Village, Pracimantoro, Wonogiri, Indonesia. Open Environ Res J, 2025; 18: e25902776419140. http://dx.doi.org/10.2174/0125902776419140251028065415



Received: June 24, 2025 Revised: July 25, 2025 Accepted: August 01, 2025 Published: November 12, 2025



Send Orders for Reprints to reprints@benthamscience.net

1. INTRODUCTION

The very low carrying capacity of the karst area makes it prone to dissolution because it is vulnerable to climate change. Indonesia has a karst area of 15.4 million hectares, or 20% of the total area of Indonesia (Apriliana & Budiyanto, 2019). The rocks that make up the karst area contain carbonate rocks, making them easily damaged, especially when they come into contact with water with an acidic pH. Karst is a unique ecosystem because of the

hydrological cycle below the surface of the soil, which means that only certain vegetation with low water requirements can grow in the karst area (Prabudimas & Budianto, 2020). This is because surface water flows into the holes in the cave system to form underground rivers (Faida *et al.*, 2018). The underground water flow in the karst area in Indonesia, particularly in Pracimantoro, had a water discharge of 7,900 liters/second with 76 springs in 1998–1999 (Samodra, 2005).

The development of community forests and limited agriculture in the karst of Pracimantoro District is carried out as a form of area preservation. Hudiyani *et al.* (2017) stated that forests are one of the pillars of community life, maintain ecosystem balance, and provide a means for social interaction. According to Salim and Pratiwi (2015), the level of plant diversity in an area affects soil fertility. The circulation of nutrients in tropical forest areas occurs specifically where vegetation functions as a provider of nutrient reserves. Most of the nutrients in the forest are stored in forest stands, and some are in the soil layer (Siagian *et al.*, 2021). The amount of nutrient input to an area is influenced by litter from the vegetation above it through the decomposition process (Tongkaemkaew *et al.*, 2018).

Forest productivity depends on the amount of litter that falls, and dense forest canopies tend to contribute more litter than sparse canopies (Kusmana *et al.*, 2021). Nursal *et al.* (2015) reported that dense forest vegetation can influence the decomposition process of litter on the forest floor. A closed canopy affects the entry of light to the forest floor. Less light entering the forest floor creates more humid conditions, favoring decomposers (Wibowo & Rizqiyah, 2014). Furthermore, Wibowo & Rizqiyah (2014) stated that the increase in decomposers is directly proportional to the speed of litter decomposition.

Vegetation as a litter producer plays an important role in the forest nutrient cycle, maintaining and restoring nutrients and soil fertility. Litter that falls on the forest floor becomes a habitat for microorganisms and fungi, facilitating the decomposition process. The ability to decompose litter is influenced by environmental factors and litter chemistry (Aerts, 1997). The karst landscape is an interesting research location due to its unique ecosystem. Research on karst areas is still rare in Indonesia, including studies measuring the rate of litter decomposition. Pracimantoro District has potential community forests that have never been studied regarding litter decomposition rates. Variation in vegetation structure, species composition, and planting patterns in community forests is believed to influence microclimate the presence of decomposing conditions and microorganisms. Therefore, this research is essential to obtain information on litter decomposition rates to maintain the stability of mineral nutrients in the karst ecosystem and support the sustainability of the karst area.

2. MATERIALS AND METHODS

2.1. Site Study and Research Design

The study was conducted in the Pracimantoro karst ecosystem community forest, Wonogiri Regency. The study was carried out in community forests with three different types of land management: monoculture, polyculture, and agroforestry. This study aims to determine the rate of decomposition in the three community forest planting patterns.

This study is an analytical study conducted with a quantitative approach, focusing on comparing the

percentage of litter mass loss from each land area, with environmental parameters used as control variables. The study was conducted observationally by placing litter bags on each land-use pattern and observing them for a specified period without any manipulation.

Measurement plots were determined by purposive sampling, considering the representativeness of land management patterns (Tongkaemkaew $et\ al.$, 2018). Field data collection from each type of community forest land management was carried out by creating plots with a 3-case \times 3 system \times 3-repetitions design, resulting in a total of 9 measurement plots (Tongkaemkaew $et\ al.$, 2018). The data collection plots were square with an area of 0.04 ha, as commonly used in karst areas (Salim & Budiadi, 2014; Moro $et\ al.$, 2016; Tongkaemkaew $et\ al.$, 2018). Furthermore, three points intersecting the diagonal of each measurement plot were selected as locations for placing research samples, with 12 litter bags placed in each plot (Tongkaemkaew $et\ al.$, 2018).

2.2. Literature Review Strategy

A literature review was conducted to support the interpretation of field data and to contextualize the decomposition rate of leaf litter in karst ecosystems. The literature search was carried out using several academic databases, including ScienceDirect, Scopus, Google Scholar, and DOAJ, between January and April 2024.

This study involved the collection of leaf litter from cultivated tree species commonly found in community forests (e.g., Swietenia mahagoni, Tectona grandis, and Acacia auriculiformis). All plant materials were collected non-destructively from the forest floor and did not involve endangered or protected species as listed by the IUCN Red List or CITES. The research complied with all relevant institutional, national, and international guidelines on plant research. Fieldwork was conducted with the permission of local authorities and with the cooperation of forest farmer groups in Wonodadi Village, Pracimantoro, Wonogiri, Indonesia.

2.3. Data Collection

2.3.1. Litter decomposition

The measurement of the litter decomposition rate was carried out using the litter bag method. Data were collected with a litter bag weighing 20 grams/bag by arranging the litter, especially on large leaves, so that layering does not occur, thus slowing down microbial respiration in decomposition (Berg & McClaugherty, 2008; Devianti & Tjahjaningrum, 2017). And then, a litter bag was placed on the forest floor (Devianti & Tjahjaningrum, 2017; Susanto et al., 2021) and taken every two weeks (Salim & Budiadi, 2014; Karina et al., 2022). Furthermore, the litter will be dried using an oven at a temperature of 105 °C for 24 hours until the weight is constant (Kusmana & Yentiana, 2021), then the weight reduction is calculated. The analysis of the decomposition rate uses the Olson (1963) equation as shown in Eq. (1) (Aerts, 1997; Kusmana & Yentiana, 2021). Data analysis was carried out using SPSS version 16.

$$Xt = X0.e^{-kt} \tag{1}$$

Notes: k = litter decomposition rate; T = observation time (weeks)

2.3.2. Environmental Parameters

Environmental parameter measurements were conducted every 15 days in the morning (07:00 - 09:00 WIB), afternoon (12:00 - 14:00 WIB), and evening (16:00 - 18:00 WIB) (Annisa et al., 2015) along with the day the samples were taken. Measurement of temperature and humidity using a thermohygrometer. The results obtained were then presented as a daily relative average table. Data analysis was calculated using Eq. (2) (Setyowati, 2008; Annisa et al., 2015; Simbolon et al., 2022).

$$E = \frac{(2 \times E1) + E2 + E3}{4} \tag{2}$$

Notes: E = environmental parameters; E1 = morning measurement; E2 = afternoon measurement; E3 = evening measurement

3. RESULT AND DISCUSSION

3.1. Vegetation of the Karst Community Forest

The results of vegetation research in the Pracimantoro karst community forest are presented in Table 1. Pracimantoro District is a karst mountain area whose main structure is limestone and carbonate rock. The development of community forests in karst areas was chosen to rehabilitate land and preserve the hydrological system (Nasrudin & Parikesit, 2020). Community forest management is carried out traditionally by implementing three forms of planting patterns, namely monoculture (Fig. 1a), polyculture (Fig. 1b), and agroforestry community forests (Fig. 1c). In addition to forestry plants, several species of understory plants were also found as medicinal plants and crops, especially in agroforestry patterns. The

medicinal plants found were turmeric (*Curcuma longa*), lempuyang (*Zingiber zerumbet*), and ginger (*Zingiber officinale*), while taro (*Colocasia esculenta*) is a crop.

Table 1. Type of pracimantoro karst community forest management based on planting patterns.

Community Forest Planting Patterns	Tree Species	Other Components		
Agroforestry	Teak, Sengon, Acacia, Sandalwood, Mahogany, Melinjo, Petai, Rosewood	<u>Livestock:</u> Cows, Bees <u>Agriculture:</u> Ginger, Turmeric, Taro, Lempuyang		
Polyculture	Acacia, Mahogany, Teak	=		
Monoculture	Mahogany, Acacia, Bintaro	-		

The results of species identification in the three forms of community forest management (Fig. 1 and Table 1) are divided into six families: Fabaceae, Meliaceae, Lamiaceae, Gnetaceae, Santalaceae, and Apocynaceae. According to Prabowo et al. (2024), the stands originating from the Fabaceae and Meliaceae families were the dominant ones. Previously, Faida et al. (2018) also reported that the karst ecosystem is composed of vegetation with long and deep roots, such as Tectona grandis, Swietenia sp., Acacia auriculiformis, Paraserianthes falcataria, and Melaleuca cajuputi. As shown in Table 2, Mahogany (Swietenia mahagoni), Teak (Tectona grandis), and Acacia (Acacia auriculiformis) were the most dominant species and had the highest IVI in the Pracimantoro karst community forest. In accordance with research conducted by Adiputra et al. (2022), Mahogany (Swietenia mahagoni) dominates the forest in the Gunung Sewu Karst area, followed by Teak (Tectona grandis). These species are trees with commercial purposes as sawmill timber (Table 2) (Nasreldin et al., 2024).




Fig. (1). Community forest management in Pracimantoro District based on planting patterns. a) monoculture, b) polyculture, and c) agroforestry.

They are selected in community forest management because they have high economic value for the community (Hudiyani et al., 2017). In addition, from an ecological perspective, these species have roots that can penetrate deep into the soil and possess high adaptive abilities to the climate in the karst area (Samodra, 2005). Mahogany stands have the highest IVI of 105.34% (Table 2). Berg & McClaugherty (2008) conveyed that species with the highest IVI have a greater chance in an ecosystem community to maintain their growth. In addition, there are species of stands that have IVI values that are not much different, namely acacia and teak species with values of 71.49% and 71.38%, respectively. The magnitude of the IVI values of the two species means that both have positions, roles, and growth abilities that are not much different in their communities (Adiputra et al, 2022). These results show that these species have a higher position than other species found, especially their ability to adapt to the environment as indicated by the magnitude of the IVI value (Pertiwi & Chairul, 2024).

Vegetation plays an important role in the formation of community forests. Based on Table 2, it can be concluded that the community forest in Wonodadi Village has a dense forest with a stand density of 5,275 ind/ha. Furthermore, the highest density is given by mahogany, teak, and acacia stands (Table 2). High density illustrates that the species has the largest number of other species at that location (Zulkarnain *et al.*, 2015). This is then in accordance with the condition of the vegetation planted, without taking into account the planting distance and without carrying out thinning activities.

3.2. Environmental Parameters

3.2.1. Temperature

The environmental parameters at the study site are presented in Table 3. The average daily air temperature in the Pracimantoro karst community forest is 30.5°C. This value is slightly higher than the average air temperature

in karst ecosystem forests in China, which was 30.3° C (Zhu *et al.*, 2021). Based on the classification index by Setyowati (2008), the average air temperature in the Pracimantoro karst community forest indicates that the area has a hot microclimate.

Community forests with a monoculture pattern have the highest average daily temperature of 31.65°C. The high temperature is due to the species planted on monoculture land having light canopies. The thin canopy allows high light intensity to penetrate to the forest floor, resulting in low air humidity under the stand (Siagian *et al.*, 2021). Indirectly, air temperature significantly affects the reduction of litter mass because it also influences microbial activity (Deviani & Tjahjaningrum, 2017).

3.2.2. Relative humidity

Based on Table 3, the average relative humidity in the karst community forest was 62.55%, with the highest average of 64.82% in agroforestry land. Susanti & Halwany (2017) stated that air temperature in an area is inversely proportional to its air humidity. This is consistent with the results of this study, showing that community forests with higher humidity have relatively lower air temperatures. The classification of climate conditions based on air humidity by Setyowati (2008) indicates that the Pracimantoro karst community forest has a dry climate, as it has a relative humidity of <70%.

Air humidity is a critical environmental factor in the litter decomposition process because it affects the presence of decomposing microorganisms. Appropriate humidity (>70%) can accelerate litter decomposition even in areas with high air temperature (Deviani & Tjahjaningrum, 2017). Jupri et al. (2024) stated that air humidity in an area is influenced by air temperature, light intensity, air pressure, water availability, and vegetation. Agroforestry community forests have vegetation with medium to dense canopies, creating higher air humidity. However, the karst community forest, with air humidity below 70%, experiences a slower rate of decomposition.

Table 2. Important values indeks (IVI) of tree species in the pracimantoro karst community forest.

No	Species	Family	K (n/ha)	KR (%)	FR (%)	DR (%)	IVI (%)
1	Swietenia mahagoni	Meliaceae	2150	23.70	24.00	40.58	105.34
2	Acacia auriculiformis	Fabaceae	1125	40.76	24.00	26.16	71.49
3	Tectona grandis	Lamiaceae	1250	21.33	24.00	23.68	71.38
4	Cerbera manghas	Apocynaceae	400	7.58	4.00	3.03	14.61
5	Paraseriantes falcataria	Fabaceae	125	1.90	4.00	2.24	8.61
6	Gnetum gnemon	Gnetaceae	100	2.37	4.00	1.51	7.40
7	Parkia speciosa	Fabaceae	25	0.47	4.00	1.23	5.70
8	Santalum album	Santalaceae	50	0.95	4.00	0.38	5.33
9	Swietenia macrophylla	Meliaceae	25	0.47	4.00	0.79	5.27
10	Dalbergia latifolia	Fabaceae	25	0.47	4.00	0.38	4.86
Total			5275	100	100	100	300

Polyculture

Monoculture

Average

87.55

83.53

83.06

-0-000				
Community Forest Planting Patterns	Temperature (°C)	Relative humidity (%)	Soil pH	Light Intensity (%)
Agroforestry	29.96	64.83	7,3 - 7,8	78.11

63.98

58.87

62.55

29.88

31.65

30.50

Table 3. The environmental parameters in different planting patterns of the Pracimantoro karst community forest.

3.2.3. Soil pH

Based on the results of field analysis, the soil pH in the Pracimantoro karst community forest ranged from 7.0 to 8.0. The research by Prabowo et al. (2024) also showed that the soil pH in the Pracimantoro karst community forest ranged from 7.56 to 8.03. This indicates that the soil in the karst community forest is neutral to slightly alkaline. The karst structure, consisting of limestone (CaCO₃), results in high base cations, making the soil alkaline. Muhsin et al. (2017) stated that the optimal soil pH for litter decomposition is 6.5 to 7.0. Furthermore, Nursal et al. (2015) added that alkaline soil can interfere with the activity of decomposing microorganisms, causing the decomposition process to proceed slowly. In their research, Susanto et al. (2021) concluded that soil pH has a 50% influence on the litter decomposition process when combined with appropriate humidity.

3.2.4. Rainfall

Rainfall data obtained from the Central Statistics Agency of Wonogiri Regency over a period of 10 years (2014-2023) show that Wonogiri Regency has rainfall ranging from 1,108 to 3,066mm/year, with an average of 2,084.1mm/year. According to the classification by Schmidt and Ferguson, Wonogiri Regency falls under climate type C (rather wet), with a Q value of 55.89%.

Based on data from the Ministry of Public Works and Spatial Planning, Directorate General of Water Resources, BBWS Bengawan Solo (2024), rainfall in Wonogiri Regency averaged 134.43mm, and during observations from May to July, it averaged 25.89mm. According to the Meteorology, Climatology, and Geophysics Agency (BMKG), the average rainfall in that year indicates that Wonogiri Regency experiences medium rainfall, while the average monthly rainfall during that period is classified as low. Deviani & Tjahjaningrum (2017) stated that rainfall affects humidity and environmental temperature, and high rainfall can facilitate litter breakdown, accelerating decomposition. The low rainfall during the research period made temperature and humidity less supportive of the decomposition process.

3.2.5. Light Intensity

The light intensity in the three types of planting patterns in the community forest ranged from 78.11% to

87.55%, with an average of 83.06% (Table 3). Light intensity in different types of community forests is influenced by canopy density. Siagian *et al.* (2021) stated that a dense canopy creates shade, preventing sunlight from reaching the forest floor. The light intensity produced by solar radiation can also accelerate the photodegradation of recalcitrant chemical compounds (Huang *et al.*, 2024).

7,5 - 8

7 -7,8

7 - 8

Community forests with an agroforestry planting pattern have the lowest light intensity. The diverse strata and wide canopy of the constituent stands are influencing factors (Siagian $et\ al.$, 2021). In contrast, monoculture land has the highest light intensity because the planting pattern includes Acacia and Bintaro, both of which have light canopies. The temperature and humidity of the air beneath the stand are also impacted by light intensity (Nurjanto $et\ al.$, 2016). Denser stands block more sunlight from reaching the forest floor, resulting in lower light reception (Simbolon $et\ al.$, 2022), which in turn increases humidity levels.

3.3. Litter Decomposition Rate

3.3.1. Litter Weight Reduction

The process of leaf litter decomposition can be seen directly by observing the color changes of the leaves along with changes in their physical structure (Fig. 2). The initial stage of litter decomposition is the occurrence of physical changes in the litter, which will become more intensive over time (Karina et al., 2022). Color changes in leaf litter can occur due to the breakdown of the green leaf substance (chlorophyll) in the leaves, which were originally green, turning yellow, brown, and changing to other darker colors (Berg & McClaugherty, 2008). Physical changes in leaf litter during observation did not show any striking changes (Fig. 2). However, the structure of the litter began to become brittle and break and crack. Changes in litter structure are presented in Fig. (2a) (monoculture), Fig. (2b) (polyculture), and Fig. (2c) (agroforestry). Damage to the leaf litter structure on monoculture and agroforestry is more visible when compared to damage to the leaf litter structure on polyculture land. Siagian et al. (2021) stated that the decomposition process will change the structure of leaf litter to become more brittle and then form flakes.




Fig. (2). Physical changes of litter in karst community forests for 60 days. (a) Monoculture; (b) Polyculture; (c) Agroforestry.

Dominantly, the litter decomposition process is accompanied by a decrease in the dry weight of the litter. The diverse composition of the stand causes variation in litter weight loss rates in the field. Litter weight loss occurs fluctuatingly depending on supporting factors over time (Nursal $et\ al.$, 2015). The decrease in litter weight in the Pracimantoro karst community forest is presented in Fig. (3).

The average final litter weight, based on Fig. (3), in agroforestry, polyculture, and monoculture community forests was 13.04 g, 14.23 g, and 12.90 g, respectively. The most significant decrease in litter weight was observed in monoculture community forests, at 35.50%. This value is slightly higher than the weight loss in agroforestry community forests, which reached only 34.78% of the initial weight. The magnitude of litter weight loss can be

influenced by various factors, including environmental and climatic conditions, as well as the quality of the litter from the standing species.

The results shown in Fig. (4) indicate that the average percentage of dry litter weight loss in all types of community forests over 2 months was 33.05%. The

decrease in dry litter weight increased over time. The percentage decrease in litter weight observed in this study is smaller than that reported by Rumambi *et al.* (2018), who found a weight loss of 32.37% during one month of decomposition, but larger than that reported by Karina *et al.* (2022), who observed a 10.4% decrease over 3 months.




Fig. (3). Graph of litter weight reduction in Pracimantoro Karst Community Forest.




Fig. (4). Average percentage of litter weight reduction in Pracimantoro Karst community forest.

Consider	Chemical Content (%)					C	
Species	P	N	С	Lignin	Polyphenols	Sources	
Tectona grandis	0.08	0.93	48.7	15.18	11.85	Umayya, 2017	
Swietenia sp.	0.48	0.82	18.1	16.6	25.26	Umayya, 2017	
Acacia auriculiformis	0.29	3.9	49.6	19.6	7.7	Oyun et al., 2006	

Table 4. Chemical content of litter from dominant species in Pracimantoro karst community forest.

Polyculture community forests with environmental conditions that are almost similar to agroforestry forests tend to experience the slowest decrease in litter weight (Fig. 4). The decrease in litter weight is only 28.85% of its initial weight (Fig. 3). This is due to the dominance of stands with low litter quality so that they are difficult to decompose. These stands include: mahogany, teak, and acacia. Susanto $et\ al.\ (2021)$ stated that the loss of litter weight is caused by differences in litter chemistry.

Based on Table 4, the three dominant species in the study site have high lignin and phenol chemical content, so they have low litter quality and are difficult to decompose. Salim & Budiadi (2014) emphasized that a land will experience a slow decomposition process if it is dominated by stands that have high lignin and polyphenol content in their litter. According to Hairiah et al. (2006), high-quality litter is litter with a C/N ratio of <25%, lignin <15%, and polyphenol content <3% which will be easily decomposed. Lignin and polyphenol content are difficult to decompose because of their resistance to decomposer bacteria (Darmayanti & Rindyastuti, 2012). Aerts (1997) stated that the lignin content of evergreen stands is relatively high, so that it can control the speed or slowness of the litter decomposition process. Mahogany and Acacia have higher lignin than teak because both are included in evergreen stands, while teak is not (Table 4).

In high concentrations, lignin and polyphenol content can inhibit the rate of decomposition, whereas the N and P $\,$

content in the litter can accelerate the rate of decomposition (Aerts, 1997; Huang et al., 2024). Therefore, the litter weight loss in monoculture community forests was highest because mahogany and acacia, as their constituent species, have higher N and P content than teak, which is not their constituent species (Table 4). Furthermore, agroforestry community forests have the second-highest weight loss below monoculture forests because, in the agroforestry land, the three stands above are not the primary constituent stands but are combined with other stands.

3.3.2. Litter Decomposition Rate

The rate of leaf litter decomposition in several forms of community forest land management is presented in Fig. (5). The litter decomposition rate was illustrated by the decomposition constant (k). The results of the analysis show that the average decomposition rate is 0.0268. The highest decomposition rate is on land with a monoculture planting pattern of 0.0293, then in an agroforestry planting pattern of 0.0277, and the smallest decomposition rate of 0.0229 in community forests with a polyculture planting pattern (Fig. 5). The decomposition rate is influenced by environmental conditions such as vegetation, temperature, air humidity, and rainfall. [16] Vegetation density and the amount of rainfall are the main factors causing high or low temperatures and air humidity in an area.




Fig. (5). The litter decomposition rate (k) in several planting patterns of community forest management in the Pracimantoro Karst ecosystem. The X-axis represents the type of planting pattern, while the Y-axis shows the decomposition rate constant (k) as calculated from dry mass loss.

The decomposition rate was calculated from the weight loss that occurred during the research process. The litter decomposition process occurs in two stages, namely the reduction in water content of the litter along with dissolved chemicals in water, such as glucose and phenol compounds, and further decomposition with the breakdown of complex litter chemicals such as lignin and cellulose (Nursal *et al.*, 2015).

Based on the measurements carried out, Wonodadi Village has low rainfall, a dry and hot climate, so the decomposition process is slow. It was recorded that the Pracimantoro karst community forest had a very low k value when compared to research at the Nglanggeran Gunungkidul Ancient Volcano by Moro et al. (2016), which had a positive k value ≤ 3 as a weathering constant. However, the results of the measurement of the litter decomposition rate are higher than the research of Karina et al. (2022) at KHDTK, Bengkulu University, whose highest decomposition constant value was <0.02. Even so, the decomposition rate that occurs in the Pracimantoro karst community forest is difficult to observe because it has a very small k value (<0.1). Moro et al. (2016) stated that the decomposition constant value (k) <0.1 will be difficult to observe, especially during the dry season.

Based on Fig. (5), the decomposition rate that occurs in the karst community forest has a number that is not much different from one another. The k value tends to increase in all forms of land management as the observation time increases. Deviani & Tjahjaningrum

(2017) stated that the suppression effect of litter decomposition will occur since the litter shrinks by 30 - 40% of the initial weight of organic material. Microclimate conditions in an area will affect the presence of decomposing microorganisms and affect their speed in decomposing litter (Karina *et al.*, 2022).

An environment that is always wet and humid will accelerate the decomposition process of litter on the forest floor. The hot and dry environmental conditions in the Pracimantoro karst forest have an impact on the slow rate of decomposition in the area. Berg & McClaugherty (2008) said that the value of the litter decomposition rate can be interpreted as the rate of return of nutrients in an ecosystem. The rate of return of soil nutrients depends on the rate of litter destruction and the decrease in dry weight of litter over a certain period of time.

3.4. Model of the Relationship between dry Litter Weight and Time

The relationship model of dry weight of the remaining litter with time can be described using an exponential regression model (Fig. 6). The rate of litter decomposition is a dynamic process. Litter decomposition has different dimensions over time because it is highly dependent on factors that influence the rate of litter decomposition (Deviani & Tjahjaningrum, 2017). The longer the decomposition process takes place, the litter will continue to experience a decrease in litter weight until the litter reaches the final stage of composting.




Fig. (6). Exponential regression model of the relationship between remaining litter dry weight and decomposition time (weeks). The X-axis represents the duration of decomposition (weeks), while the Y-axis shows the remaining litter mass (grams). The curve illustrates the exponential decay trend of litter mass over time.




Fig. (7). Estimation of leaf litter weight in several types of community forest management in the Pracimantoro Karst ecosystem based on Olson's exponential regression model. The X-axis represents time (weeks), and the Y-axis represents the estimated remaining dry litter weight (grams). Shorter half-life values indicate faster decomposition rates.

Based on the exponential equation created, the decomposition process in the litter in the Pracimantoro karst community forest cannot be predicted when it will reach zero. However, the reduction in the weight of the remaining litter can be estimated using half-life calculations. Siagian et al. (2021) said that an R-square value of more than 0.5 indicates a good relationship between the two variables. The decay of the remaining weight of the litter is suitable to be presented in an exponential graph because the initial process of weight loss will be greater and slow down over time (Mahmudi et al., 2011). Nurianto et al. (2016) stated that the magnitude of the decrease in litter weight is never constant but always changes over time. From the equations obtained from the three forms of community forest management, it can be estimated that the cycle of decreasing the remaining weight in the litter decomposition process from monoculture forests, polyculture forests, and agroforestry sequentially occurs every 23.64 weeks, 30.52 weeks, and 24.31 weeks (Fig. 7). The smaller the half-life value, the faster the decomposition process will take place. Zhu et al. (2021) stated that the average litter in tropical forests throughout the world will decompose for 0.50 - 3.03 years. leaf litter experienced variations However, decomposition time between 1 - 5 years (Prescott, 2010).

The length of the litter decomposition process in the Pracimantoro karst community forest shows varying results, with a notable difference among the three community forest land management types. Based on its half-life, leaf litter in agroforestry and monoculture lands

has a similar decomposition time. The rate of litter decomposition is influenced by the type of stand, environmental conditions, and litter quality (Aerts, 1997; Prescott, 2010; Huang et al., 2024). Environmental conditions that are suitable continuously and over a long period have a significant impact on the rate of litter decomposition (Prescott, 2010; Rumambi et al., 2018). However, Prescott (2010) emphasized the importance of including a moisture threshold in Olson's decomposition model, considering that karst areas are dry and experience water limitations. This modification provides a more realistic description of decomposition dynamics in dry karst areas.

4. STUDY LIMITATIONS

This study has several limitations. First, the limited observation time may not reflect the entire litter decomposition cycle, which can last for 1-5 years depending on the type of litter and microclimate conditions. Second, the dry season during data collection caused high temperatures and low humidity, which may slow down the activity of microorganisms and affect the accuracy of the decomposition rate estimation. Third, this study only observed macro-environmental variables (temperature, humidity, pH, and light intensity) and did not measure the diversity of soil microbes or fauna, which also play an important role in the decomposition process. In the future, further research should be conducted during the rainy season and involve further biological parameters and chemical tests of litter and soil. In addition,

considering that the monoculture pattern has the fastest estimated decomposition time compared to other planting patterns, the suggestion for further research is to conduct a study on the influence of seasons (multipool) and interactions between litter in polyculture and agroforestry planting patterns on the rate of litter decomposition.

CONCLUSION

The average decrease in litter weight across all forms of karst community forest management was 33.05%. Monoculture forests had the highest decrease in litter weight at 35.50%, followed by agroforestry forests at 34.78%, and polyculture forests at 28.82%. The dominance of Mahogany (Swietenia sp.), Acacia (Acacia auriculiformis), and Teak (Tectona grandis) as constituent species greatly influenced the rate of litter decomposition, as these three species have high lignin and polyphenol content. The average rate of leaf litter decomposition in karst community forests was 0.0268. The half-life of litter decay in monoculture, polyculture, and agroforestry lands was 23.64 weeks, 30.52 weeks, and 24.31 weeks, respectively.

AUTHOR'S CONTRIBUTIONS

S.R.R., N.M.: Conception and design; S.R.R.: Data collection; S.R.R., N. M., S.: Analysis and interpretation of results; S.R.R., N.M., S.: Draft manuscript. All authors reviewed the results and approved the final version of the manuscript.

LIST OF ABBREVIATIONS

IVI = Important Value Index

k = k-value = constant decomposition rate

P = Phosphorus

N = Nitrogen

C = Carbon

CONSENT FOR PUBLICATION

Not applicable.

AVAILABILITY OF DATA AND MATERIALS

All data generated or analyzed during this study are included in this published article.

FUNDING

None.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

Thanks to the Pracimantoro District Government for granting permission to conduct this research and to the Extension Officer of the Forestry Service Branch Office Region XI for assisting with the research. Thanks also to all members of the forest farmer groups in Wonodadi, who were cooperative and facilitated research activities in the field, allowing the necessary data to be collected.

REFERENCES

- Adiputra, SW (2024) study of the structure and composition of karst ecosystem vegetation in kemadang village. Thesis, Doctoral dissertation, STIPER Yogyakarta Agricultural Institute.
- Aerts, R (1997) Climate, leaf litter chemistry and leaf litter decomposition in terrestrial ecosystems: A triangular relationship. Oikos, 79, 439-49. [http://dx.doi.org/10.2307/3546886]
- Annisa, N, Kurnain, A, Indrayatie, ER & Peran, SB (2015) Microclimate and discomfort index of city parks in komet village. *EnviroScienteae*, 11, 143-51. [http://dx.doi.org/10.20527/es.v11i3.1104]
- Apriliana, T & Budiyanto, E (2019) Dynamics of land cover and use in the karst spring catchment area of goa gremeng, ponjong district, Gunungkidul Regency. Swara Bhumi E-Journal, 1, 1-9.
- Berg, B (2008) Plant Litter: Decomposition, Humus Formation, Carbon Sequestration. Springer, Cham.
- Darmayanti, AS & Ridesti, R (2012) Changes in the percentage of litter nutrients due to the decomposition process in four species of deciduous plants in the purwodadi botanical garden. *Indo J Biol*, 8, 45-55.
- Devianti, OKA & Tjahjaningrum, ITD (2017) Study of litter decomposition rate in pine forests in the Taman Safari Indonesia II Tourism Area, East Java. ITS J Sci Art, 6, 105-9.

[http://dx.doi.org/10.12962/j23373520.v6i2.27535]

- Faida, LRW (2018) Gunung Sewu: Uncovering the traces of flora history, reconstructing the karst area. UGM Press, Yogyakarta.
- Hairiah, K, Sulistyani, H, Suprayogo, D, Widianto, P, Purnomosidhi, P, Widodo, RH & Van Noordwijk, M (2006) Litter layer residence time in forest and coffee agroforestry systems in Sumberjaya, West Lampung. For Ecol Manage, 224, 45-57.

[http://dx.doi.org/10.1016/j.foreco.2005.12.007]

- Huang, H, Wang, H, Sun, L, Mo, Z, Zhang, Y & Wei, S (2024) Effects of drought and litter types on litter decomposition in desert-oasis ecotone. Appl Soil Ecol, 204, 105742. [http://dx.doi.org/10.1016/j.apsoil.2024.105742]
- Hudiyani, I, Purnaningsih, N, Asngari, PS & Hardjanto, H (2017)
 Farmers' Perceptions of community forests with agroforestry patterns in wonogiri regency, central java province. *Extension J*, 13, 64-78.

 [http://dx.doi.org/10.25015/penyuluhan.v13i1.14709]
- Jupri, A, Ahyadi, H, Uzma, S, Muthma'innah, E, Riski, TNA, Hakim, A, Wulandari, SY & Hidayah, N (2024) Analysis of the effect of climate change on biodiversity conditions in West Nusa Tenggara. *Jurnal Biologi Tropis*, 24, 1-13. [http://dx.doi.org/10.29303/jbt.v24i2b.7941]
- Karina, TP, Arianto, W & Wiryono, W (2022) Leaf litter decomposition rate in special purpose forest area (KHDTK) of Bengkulu University, North Bengkulu. J Global ForestEnvironment Sci, 2, 106-12.
- Kusmana, C & Yentiana, RA (2021) Leaf litter decomposition rate of *Shorea guiso* in Dramaga Research Forest, Bogor, West Java. *Journal Of Tropical Silviculture*, 12, 172-7. [http://dx.doi.org/10.29244/j-siltrop.12.3.172-177]

- Mahmudi, M. Soemarno, M & Arfiati, D (2011) Production and decomposition of rhizophora mucronata litter and its contribution to nutrients in reforestation Mangrove Forest, Nguling Pasuruan. I Berkala Penelitian Hayati, 6, 19-24.
- Moro, HKEP, Zulfikar, M, Wibowo, MS & Recto, S (2016) Leaf Litter Decomposition Rate on the Forest Floor of Nglanggeran Ancient Volcano. Proceedings of the National Seminar on Wallacea Biodiversity.
- Muhsin, Ahmad Sw (2017) Decomposition rate of Ironwood Leaf Litter (Metrosideros petiolata Coord.) in the Nanga-Nanga Papalia protected forest Area, Kendari City, Southeast Sulawesi. Biowallacea, 4, 655-66.
- Gurashi, NA, Yasin, EHE & Czimber, K (2024) Assessment of tree species availability based on sawmilling and Timber Markets Survey in Sinnar State, Sudan. Acta Silv Lignaria Hung, 20, 39-51. [http://dx.doi.org/10.37045/aslh-2024-0003]
- Nasrudin, A (2020) Vegetation Analysis Of Karst In Padjadjaran University Campus Area Cintaratu, Pangandaran, West Java. Proceedings of the National Seminar of the Indonesian Biodiversity Society.
- Nurjanto, H H, Supriyo, H, Widyastuti, S M & Kabirun, S (2016) Decomposition of various types of gamal litter in the wanagama 1 educational forest, gunung kidul, yogyakarta. Tropical Forestry J, 6137
- Nursal, N, Syafi'i, WSI & Hanif, MAHA (2015) The rate of leaf litter decomposition in the rumbio customary forest area, kampar district, kampar regency as a development of learning modules on the concept of tropical forest ecosystems in class x high school. Thesis, Doctoral Dissertation, Riau University.
- Oyun, MB, Kadeba, O & Aletor, VA (2006) Nitrogen release patterns of mixed Gliricidia sepium and Acacia auriculiformis leaves as influenced by polyphenol, lignin and nitrogen contents. J Applied Sci, 6, 2217-23. [http://dx.doi.org/10.3923/jas.2006.2217.2223]
- Pertiwi, V & Chairul, C (2024) Analysis of tree-level vegetation in the Silokek Geopark Area, Sijunjung Regency. Sci J Biol, 12, 367-79. [http://dx.doi.org/10.33394/bioscientist.v12i1.9231]
- Prabowo, SH, Rahmadwiati, R & Nufus, M (2024) Exploring the vegetation characteristics of karst landscapes: A study of
 - community forest in Tubokarto Village, Wonogiri, Indonesia. Media Konservasi, 29, 593-3.
 - [http://dx.doi.org/10.29244/medkon.29.4.593]
- Prabudimas, CA & Budiyanto, E (2020) Characterization of interhill valley vegetation cover in the East Gunungsewu Karst Area in Ponjong District. Swara Bhumi, 1, 1-7.
- Prescott, CE (2010) Litter decomposition: What controls it and how can we alter it to sequester more carbon in forest soils? Biogeochemistry, 101, 133-49. [http://dx.doi.org/10.1007/s10533-010-9439-0]
- Rumambi, JF, Langi, MA & Nurmawan, W (2018) Initial decomposition rate of tree litter Palaquium obovatum,

- Spathodea campanulata and Calophyllum soulattri in Bron Warembungan Forest, Minahasa Regency. Eugenia, 24 [http://dx.doi.org/10.35791/eug.24.3.2018.22791]
- Salim, AG & Budiadi, B (2014) Production and nutrient content of litter in nglanggeran community forest, Gunung Kidul, DI Yogyakarta. J Plant Forest Res, 11, 77-88. [http://dx.doi.org/10.20886/jpht.2014.11.2.77-88]
- Samodra, H (2005) Potential of Karst Natural Resources in Gunungkidul Regency, Yogyakarta. Center for Geological Research and Development, Geological Department of Energy and Mineral Resources, Bandung.
- Setyowati, DL (2008) The micro climate and the need of green open space for the city of Semarang. J Human Environment, 15, 125-40.
- Siagian, SPS, Susatya, A & Saprinurdin, S (2021) Decomposition rate of Psychotria malayana Leaf litter in the Campus Forest of the University of Bengkulu. J Global Forest Environment Sci, 1, 1-9. [http://dx.doi.org/10.29244/j-siltrop.12.3.172-177]
- Simbolon, N A (2022) Decomposition Rate of Rhizophora apiculata Bl Leaf Litter in Mangrove Forest of Panjang Beach and Baai Island Nature Tourism Park Area, Bengkulu City. J Global Forest Environment Sci, 2, 90-6.
- Susanti, PD & Halwany, W (2017) Litter decomposition and soil macrofauna diversity in Nyawai Industrial plantation forests (Ficus variegate. Blume). J For Sci, 11, 212-23. [http://dx.doi.org/10.22146/jik.28285]
- Susanto, SA, Qayim, I & Triadiati, T (2021) Characteristics of fallow land with different ages and their effect on litter decomposition in Manokwari, West Papua. I Soil Climate, 45, 117-32.
 - [http://dx.doi.org/10.21082/jti.v45n2.2021.117-132]
- Tongkaemkaew, U, Sukkul, J, Sumkhan, N, Panklang, P, Brauman, A & Ismail, R (2018) Litterfall, litter decomposition, soil macrofauna, and nutrient contents in rubber monoculture and rubber-based agroforestry plantations. Forest and Society, 2, 138-49. [http://dx.doi.org/10.24259/fs.v2i2.4431]
- Umavva, L (2017) Effect of litter type to earthworm Pontoscolex corethrurus development. Undergraduate Universitas Brawijaya, Malang, Indonesia.
- Wibowo, C & Rizgiyah, W (2014) Diversity of soil Macrofauna on various stand types in Gunung Walat University Forest, Sukabumi, West Java. J Tropical Silviculture, 5, 43-8.
- Zhu, X, Zou, X, Lu, E, Deng, Y, Luo, Y, Chen, H & Liu, W (2021) Litterfall biomass and nutrient cycling in karst and nearby non-karst forests in tropical China: A 10-year comparison. Sci Total Environ, 758, 143619.
 - [http://dx.doi.org/10.1016/j.scitotenv.2020.143619] [PMID: 33221014]
- Zulkarnain, Alimuddin (2015) Vegetation analysis and visualization of forest vegetation profiles in the Tahura Nipa-Nipa Forest ecosystem in Mangga Dua Village, Kendari City. Ecogreen, 1, 43-54.

DISCLAIMER: The above article has been published, as is, ahead-of-print, to provide early visibility but is not the final version. Major publication processes like copyediting, proofing, typesetting and further review are still to be done and may lead to changes in the final published version, if it is eventually published. All legal disclaimers that apply to the final published article also apply to this ahead-of-print version.